

**Ex. 4.3-1.**  $T(n) = T(n-1) + n$ . Guess  $T(n) \leq c \cdot n^2$ . Then

$$\begin{aligned}T(n) &\leq c(n-1)^2 + n \\&= cn^2 - 2cn + c + n \\&= cn^2 + (1-2c)n + c \\&\leq cn^2\end{aligned}$$

The last step holds as long as

$$(1-2c)n + c \leq 0$$

$$(2c-1)n \geq c$$

$$n \geq \frac{c}{2c-1}$$

The recurrence holds so long as  $c > \frac{1}{2}$  and  $n_0 > \frac{c}{2c-1}$ .

**4.3-2.**  $T(n) = T(\lceil \frac{n}{2} \rceil) + 1$ . First attempt. Guess  $T(n) \leq c \lg n$

$$\begin{aligned} T(n) &\leq c \lg \lceil \frac{n}{2} \rceil + 1 \\ &\leq c \lg(\frac{n}{2} + \frac{1}{2}) + 1 \\ &= c \lg(\frac{n+1}{2}) + 1 \\ &= c \lg(n+1) - c + 1 \end{aligned}$$

We would need this to be less than  $c \lg n \dots$

**4.3-2.**  $T(n) = T(\lceil \frac{n}{2} \rceil) + 1$ . Try again. This time, guess  $T(n) \leq c \lg(n - b)$ .

$$\begin{aligned} T(n) &\leq c \lg(\lceil \frac{n}{2} \rceil - b) + 1 \\ &\leq c \lg(\frac{n}{2} + \frac{1}{2} - b) + 1 \\ &= c \lg(n + 1 - 2b) - c + 1 \\ &\leq c \lg(n - b) \end{aligned}$$

The last part holds if  $n + 1 - 2b \leq n - b$ , so  $b \geq 1$ ; and if  $-c + 1 \leq 0$ , so  $c \geq 1$ .

**4.3-6.**  $T(n) = 2T(\lfloor \frac{n}{2} \rfloor + 17) + n$ . Guess  $cn \lg n$ . Then

$$\begin{aligned} T(n) &= 2T(\lfloor \frac{n}{2} \rfloor + 17) + n \\ &\leq 2c(\lfloor \frac{n}{2} \rfloor + 17) \lg(\lfloor \frac{n}{2} \rfloor + 17) + n \\ &\leq 2c(\frac{n}{2} + 17) \lg(\frac{n}{2} + 17) + n \\ &= c(n + 34)(\lg(n + 34) - 1) + n \\ &= cn \lg(n + 34) - cn - c34 + n \end{aligned}$$

This isn't working out.

**4.3-6.**  $T(n) = 2T(\lfloor \frac{n}{2} \rfloor + 17) + n$ . Try again, this time guess  $c(n - 34) \lg(n - 34)$ .

$$\begin{aligned} T(n) &= 2T(\lfloor \frac{n}{2} \rfloor + 17) + n \\ &\leq 2c(\lfloor \frac{n}{2} \rfloor + 17 - 34) \lg(\lfloor \frac{n}{2} \rfloor + 17 - 34) + n \\ &\leq 2c(\frac{n}{2} + 17 - 34) \lg(\frac{n}{2} + 17 - 34) + n \\ &= c(n - 34) \lg(\frac{n-34}{2}) + n = c(n - 34)(\lg(n - 34) - 1) + n \\ &= c(n - 34) \lg(n - 34) - cn + 34c + n \leq c(n - 34) \lg(n - 34) \end{aligned}$$

The last step holds if  $-cn + 34c + n \leq 0$ .

$$cn - 34c \leq n$$

$$c \geq \frac{n}{n-34}$$

Notice that as  $n$  gets bigger, the ratio gets closer to 1, but will always be slightly bigger. Pick  $c = 2$ . Then we need  $2n - 68 \geq n$ , or  $n \geq 68$ .

**4.3-9.**  $T(n) = 3T(\sqrt{n}) + \lg n$ . Let  $m = \lg n$ ,  $n = 2^m$ . Then define

$$\begin{aligned} S(m) &= T(2^m) \\ &= 3T(2^{\frac{m}{2}}) + \lg 2^m \\ &= 3T(2^{\frac{m}{2}}) + m \\ &= 3S(\frac{m}{2}) + m \end{aligned}$$

What do you do with that? Guess  $cm \lg m$ .

$$\begin{aligned} &= 3c \frac{m}{2} \lg \frac{m}{2} + m \\ &= \frac{3}{2}cm \lg m - \frac{3}{2}cm + m \end{aligned}$$

This isn't working out. In fact, the complexity class is wrong.

**4.3-9.**  $T(n) = 3T(\sqrt{n}) + \lg n$ . Again, let  $m = \lg n$ ,  $n = 2^m$ , and  $S(m) = 3S(\frac{m}{2}) + m$ . Then guess  $m^{\lg 3} - \frac{m}{2}$ . (Of course.)

$$\begin{aligned} S(m) &= 3S\left(\frac{m}{2}\right) + m \\ &= 3\left(\left(\frac{m}{2}\right)^{\lg 3} - \frac{m}{2}\right) + m \\ &= 3\frac{m^{\lg 3}}{2^{\lg 3}} - \frac{3}{2}m + m \\ &= 3\frac{m^{\lg 3}}{3} + \frac{-3+2}{2}m \\ &= m^{\lg 3} - \frac{m}{2} \end{aligned}$$

So,  $S(m) = \Theta(m^{\lg 3}) = \Theta((\lg n)^{\lg 3})$ .