

## Chapter 7 in context:

- ▶ Chapter 5 Relations: Builds on proofs about sets
- ▶ Chapter 6 Self Reference: Interlude between Chapters 5 and 7, focuses on recursive thinking
- ▶ Chapter 7 Function: Builds on proofs about relations

## Chapter 7 outline:

- ▶ Introduction, function equality, and anonymous functions (**Today**)
- ▶ Image and inverse images (Wednesday)
- ▶ Function properties, composition, and applications to programming (Friday)
- ▶ Cardinality (next week Monday)
- ▶ Countability (next week Wednesday)
- ▶ Review (Monday, Apr 18)
- ▶ Test 3, on Ch 6 & 7 (Wednesday, Apr 20)

Cross out the term/concept that was **not** used in the reading for today as a way to think about functions

A kind of machine

A form of induction

A mapping between two collections

A kind of relation

For the function  $f : X \rightarrow Y$ ,  $X$  is the \_\_\_\_\_ and  $Y$  is the

\_\_\_\_\_.

function

constant


domain

codomain

first-class value

relation



input,  
raw materials,  
parameters




function

output,  
result,  
returned value


Alice	x3498
Bob	x4472
Carol	x5392
Dave	x9955
Eve	x2533
Fred	x9448
Georgia	x3684
Herb	x8401






**Not a function.**

(There's a domain element that is related to two things.)



**Not a function.**

(There's a domain element that is not related to anything.)



**A function.**

(It's OK that two domain elements are related to the same thing and one codomain element has nothing related to it.)

## Definition of function

Informal: A *function* is a relation in which everything in the first set is related to *exactly one thing* in the second set.

Formal:  $f \subseteq X \times Y$  is a *function* if

$\forall x \in X, \quad \exists y \in Y \mid (x, y) \in f$  **existence** of  $y$

$\wedge \forall y_1, y_2 \in Y, ((x, y_1), (x, y_2) \in f) \rightarrow y_1 = y_2$  **uniqueness** of  $y$

## Change of notation

Informal: A *function* is a relation in which everything in the first set is related to *exactly one thing* in the second set.

Formal (relation notation):  $f \subseteq X \times Y$  is a *function* if

$\forall x \in X, \quad \exists y \in Y \mid (x, y) \in f$  **existence of  $y$**

$\wedge \forall y_1, y_2 \in Y, ((x, y_1), (x, y_2) \in f) \rightarrow y_1 = y_2$  **uniqueness of  $y$**

Formal (function notation):  $f \subseteq X \times Y$  is a *function* if

$\forall x \in X, \quad \exists y \in Y \mid f(x) = y$  **existence of  $y$**

$\wedge \forall y_1, y_2 \in Y, (f(x) = y_1 \wedge f(x) = y_2) \rightarrow y_1 = y_2$  **uniqueness of  $y$**

We call  $X$  the *domain* and  $Y$  the *codomain* of  $f$ .



**Definition of function equality.** Let  $f, g : X \rightarrow Y$

Old definition: functions are sets.

$$f = g \text{ if } \forall f \subseteq g \wedge g \subseteq f$$

New definition: based on function notation.

$$f = g \text{ if } \forall x \in X, f(x) = g(x)$$

Function equality:  $f = g$  if  $\forall x \in X, f(x) = g(x)$

Let  $f, g : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = x \cdot (x - 1) - 6$  and  $g(x) = (x - 3)(x + 2)$ .

Prove  $f = g$ .

The old and new definitions of function equality are equivalent.

**Ex 7.2.1.**  $(\forall x \in X, f(x) = g(x))$  iff  $(f \subseteq g \wedge g \subseteq f)$ .

The old and new definitions of function equality are equivalent.

**Ex 7.2.1.**  $(\forall x \in X, f(x) = g(x))$  iff  $(f \subseteq g \wedge g \subseteq f)$ .

**Proof.** First, suppose  $\forall x \in X, f(x) = g(x)$ , that is,  $f = g$  by definition of function equality. Further suppose  $(x, y) \in f$ . By function notation,  $f(x) = y$ . By supposition and substitution,  $g(x) = y$ . By relation notation,  $(x, y) \in g$ . Finally,  $f \subseteq g$  by definition of subset.

Similarly  $f \subseteq g$ , and therefore  $f = g$  by definition of set equality.

Conversely, suppose  $f \subseteq g \wedge g \subseteq f$ , that is,  $f = g$  by definition of set equality. Further suppose  $x \in X$ .

Let  $y = f(x)$ . Note that this  $y \in Y$  must exist by definition of function. By relation notation,  $(x, y) \in f$ .

By definition of subset [or set equality],  $(x, y) \in g$ . In function notation, that is  $g(x) = y$ , and so  $f(x) = g(x)$  by substitution. Therefore  $f = g$  by definition of function equality.  $\square$

**For next time:**

*Pg 331: 7.2.(2 & 3)*

*Pg 335: 7.3.(3, 4, 8)*

*Read 7.4*

*Skim 7.5*