

## Chapter 7 outline:

- ▶ Introduction, function equality, and anonymous functions (Monday)
- ▶ Image and inverse images (Wednesday)
- ▶ Function properties, composition, and applications to programming (**Today**)
- ▶ Cardinality (next week Monday)
- ▶ Countability (next week Wednesday)
- ▶ Review (Monday, Apr 18)
- ▶ Test 3, on Ch 6 & 7 (Wednesday, Apr 20)

## Today:

- ▶ Programming: map and filter
- ▶ Definition of one-to-one and onto, plus proofs
- ▶ Inverse functions
- ▶ Definition of function composition, plus proofs


**For next time:**

*Pg 346: 7.6.(2, 3, 6)*

*Ex "7.5.(a-c)" on Schoology*


*Pg 351: 7.8.(1, 5, 6)*

*Skim 7.9*




Not a function.

(There's a domain element that is related to two things.)



Not a function.

(There's a domain element that is not related to anything.)



### **Onto** (Surjection)

Everything in the codomain is hit.

$f : X \rightarrow Y$  is onto if  $\forall y \in Y,$   
 $\exists x \in X \mid f(x) = y.$

#### **Analytic use:**

$f$  is onto.

$y \in Y.$

Hence  $\exists x \in X$  such that  $f(x) = y.$


#### **Synthetic use:**

Suppose  $y \in Y.$

$\vdots$

*(Somehow find  $x$  such that  $f(x) = y.$ )*

Therefore  $f$  is onto.



**One-to-one** (Injection)

Domain elements don't share.

$f$  is one-to-one if  $\forall x_1, x_2 \in X$ ,  
if  $f(x_1) = f(x_2)$  then  $x_1 = x_2$ .

**Analytic use:**

$f$  is one-to-one.

$f(x_1) = f(x_2)$ .

Hence  $x_1 = x_2$ .


**Synthetic use:**

Suppose  $x_1, x_2 \in X$  and  $f(x_1) = f(x_2)$ .


$\vdots$

(Somehow show  $x_1 = x_2$ .)


Therefore  $f$  is one-to-one.



Onto  
(not one-to-one)  
 $|X| \geq |Y|$




One-to-one  
(not onto)  
 $|X| \leq |Y|$



Both onto and one-to-one  
 $|X| = |Y|$


Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = \frac{x}{2}$ . Is  $f$  one-to-one? Is it onto?



Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = \frac{x}{2}$ . Is  $f$  one-to-one? Is it onto?

$f$  is one-to-one.

**Proof.** Suppose  $x_1, x_2 \in \mathbb{R}$  such that  $f(x_1) = f(x_2)$ . [Want  $x_1 = x_2$ ] Then, by how  $f$  is defined,






Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = \frac{x}{2}$ . Is  $f$  one-to-one? Is it onto?

$f$  is one-to-one.

**Proof.** Suppose  $x_1, x_2 \in \mathbb{R}$  such that  $f(x_1) = f(x_2)$ . [Want  $x_1 = x_2$ ] Then, by how  $f$  is defined,

$$\frac{x_1}{2} = \frac{x_2}{2}$$

$$x_1 = x_2$$



Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = \frac{x}{2}$ . Is  $f$  one-to-one? Is it onto?


$f$  is one-to-one.

**Proof.** Suppose  $x_1, x_2 \in \mathbb{R}$  such that  $f(x_1) = f(x_2)$ . [Want  $x_1 = x_2$ ] Then, by how  $f$  is defined,

$$\begin{aligned}\frac{x_1}{2} &= \frac{x_2}{2} \\ x_1 &= x_2\end{aligned}$$

Therefore  $f$  is one-to-one by definition.  $\square$

$f$  is onto.



Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = \frac{x}{2}$ . Is  $f$  one-to-one? Is it onto?

$f$  is one-to-one.


**Proof.** Suppose  $x_1, x_2 \in \mathbb{R}$  such that  $f(x_1) = f(x_2)$ . [Want  $x_1 = x_2$ ] Then, by how  $f$  is defined,

$$\begin{aligned}\frac{x_1}{2} &= \frac{x_2}{2} \\ x_1 &= x_2\end{aligned}$$

Therefore  $f$  is one-to-one by definition.  $\square$

$f$  is onto.

**Proof.** Suppose  $y \in \mathbb{R}$ . [Want  $x$  such that  $f(x) = y$ .]



Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = \frac{x}{2}$ . Is  $f$  one-to-one? Is it onto?

$f$  is one-to-one.

**Proof.** Suppose  $x_1, x_2 \in \mathbb{R}$  such that  $f(x_1) = f(x_2)$ . [Want  $x_1 = x_2$ ] Then, by how  $f$  is defined,

$$\begin{aligned}\frac{x_1}{2} &= \frac{x_2}{2} \\ x_1 &= x_2\end{aligned}$$

Therefore  $f$  is one-to-one by definition.  $\square$


$f$  is onto.

**Proof.** Suppose  $y \in \mathbb{R}$ . [Want  $x$  such that  $f(x) = y$ .]


Let  $x = 2y$ . Then

$$\begin{aligned}f(x) &= \frac{2y}{2} \\ &= y\end{aligned}$$


Therefore  $f$  is onto by definition  $\square$



Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = x^2$ . Is  $f$  one-to-one? Is it onto?



Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x) = x^2$ . Is  $f$  one-to-one? Is it onto?



$f$  is not one-to-one.

$$f(2) = 2^2 = 4$$

$$f(-2) = (-2)^2 = 4$$


$f$  is no onto.

Let  $y = -1$ .


$\nexists x \in \mathbb{R}$  such that  $f(x) = -1$ .

**Ex 7.6.4.** If  $A \subseteq X$  and  $f$  is one-to-one, then  $F^{-1}(F(A)) \subseteq A$ .

(Ex 7.4.9 was, Prove  $A \subseteq F^{-1}(F(A))$ , and Ex 7.4.10 was, Find a counterexample for  $A = F^{-1}(F(A))$ .)



**Ex 7.6.5.** If  $A \subseteq Y$  and  $f$  is onto, then  $A \subseteq F(F^{-1}(A))$ .





Inverse relation:  $R^{-1} = \{(y, x) \in Y \times X \mid (x, y) \in R\}$

*Since a function is a relation, a function has an inverse, but we don't know that the inverse of a function is a function.*

If  $f : X \rightarrow Y$  is a **one-to-one correspondence**, then

$$f^{-1} : Y \rightarrow X = \{(y, x) \in Y \times X \mid f(x) = y\}$$

is the *inverse function* of  $f$ .

**Theorem 7.8** *If  $f : X \rightarrow Y$  is a one-to-one correspondence, then  $f^{-1} : Y \rightarrow X$  is well defined.*

**Proof.** Suppose  $y \in Y$ . Since  $f$  is onto, there exists  $x \in X$  such that  $f(x) = y$ . Hence  $(y, x) \in f^{-1}$  or  $f^{-1}(y) = x$ .

Further suppose  $(y, x_1), (y, x_2) \in f^{-1}$  (That is, suppose that both  $f^{-1}(y) = x_1$  and  $f^{-1}(y) = x_2$ .) Then  $f(x_1) = y$  and  $f(x_2) = y$ . Since  $f$  is one-to-one,  $x_1 = x_2$ .

Therefore, by definition of function,  $f^{-1}$  is well defined.  $\square$

Relation composition: If  $R$  is a relation from  $X$  to  $Y$  and  $S$  is a relation from  $Y$  to  $Z$ , then  $S \circ R$  is the relation from  $X$  to  $Z$  defined as

$$S \circ R = \{(x, z) \in X \times Z \mid \exists y \in Y \text{ such that } (x, y) \in R \text{ and } (y, z) \in S\}$$

Function composition: If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$ , then  $g \circ f : X \rightarrow Z$  is defined as

$$g \circ f = \{(x, z) \in X \times Z \mid z = g(f(x))\}$$

**Theorem 7.9** *If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are functions, then  $g \circ f : X \rightarrow Z$  is well defined.*


**Proof.** Suppose  $x \in X$ . Since  $f$  is a function, there exists a  $y \in Y$  such that  $f(x) = y$ . Since  $g$  is a function, there exists a  $z \in Z$  such that  $g(y) = z$ . By definition of composition,  $(x, z) \in g \circ f$ , or  $g \circ f(x) = z$ .

Next suppose  $(x, z_1), (x, z_2) \in g \circ f$ , or  $g \circ f(x) = z_1$  and  $g \circ f(x) = z_2$ . By definition of composition, there exist  $y_1, y_2$  such that  $f(x) = y_1$ ,  $f(x) = y_2$ ,  $g(y_1) = z_1$ , and  $g(y_2) = z_2$ . Since  $f$  is a function,  $y_1 = y_2$ . Since  $g$  is a function,  $z_1 = z_2$ .

Therefore, by definition of function,  $g \circ f$  is well defined.  $\square$

Function composition: If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$ , then  $g \circ f : X \rightarrow Z$  is defined as

$$g \circ f = \{(x, z) \in X \times Z \mid x = g(f(x))\}$$



Let  $f(x) = 3x$


Let  $g(x) = x + 7$

Then

$$\begin{aligned} g \circ f(x) &= f(x) + 7 \\ &= 3x + 7 \end{aligned}$$

**Ex 7.8.4.** If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto, then  $g \circ f$  is onto.


**Proof.** Suppose  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto.



**Ex 7.8.4.** If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto, then  $g \circ f$  is onto.

**Proof.** Suppose  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto.


*[Now, we want to prove “onteness.” Of which function?*



**Ex 7.8.4.** If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto, then  $g \circ f$  is onto.

**Proof.** Suppose  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto.


*[Now, we want to prove “onteness.” Of which function?  $g \circ f$ . How do we prove ontteness?]*



**Ex 7.8.4.** If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto, then  $g \circ f$  is onto.

**Proof.** Suppose  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto.

*[Now, we want to prove “onteness.” Of which function?  $g \circ f$ . How do we prove ontteness? We pick something from the codomain of the function we’re proving to be onto and show that it is hit. What is the codomain of  $g \circ f$ ?*




**Ex 7.8.4.** If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto, then  $g \circ f$  is onto.

**Proof.** Suppose  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto.

*[Now, we want to prove “onteness.” Of which function?  $g \circ f$ . How do we prove ontteness? We pick something from the codomain of the function we’re proving to be onto and show that it is hit. What is the codomain of  $g \circ f$ ?  $Z$ .]*

Further suppose  $z \in Z$ . *[We need to come up with something in the domain of  $g \circ f$  that hits  $z$ . The domain is  $X$ . We will use the fact that  $f$  and  $g$  are both onto.]*






**Ex 7.8.4.** If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto, then  $g \circ f$  is onto.

**Proof.** Suppose  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are both onto.

*[Now, we want to prove “onteness.” Of which function?  $g \circ f$ . How do we prove ontteness? We pick something from the codomain of the function we’re proving to be onto and show that it is hit. What is the codomain of  $g \circ f$ ?  $Z$ .]*

Further suppose  $z \in Z$ . *[We need to come up with something in the domain of  $g \circ f$  that hits  $z$ . The domain is  $X$ . We will use the fact that  $f$  and  $g$  are both onto.]*




By definition of onto, there exists  $y \in Y$  such that  $g(y) = z$ . Similarly there exists  $x \in X$  such that  $f(x) = y$ . Now,

$$\begin{aligned} g \circ f(x) &= g(f(x)) && \text{by definition of function compos} \\ &= g(y) && \text{by substitution} \\ &= z && \text{by substitution} \end{aligned}$$

Therefore  $g \circ f$  is onto by definition.  $\square$

**Ex 7.8.5.** If  $f : X \rightarrow Y$ ,  $g : X \rightarrow Y$  and  $h : Y \rightarrow Z$ ,  $h$  is one-to-one, and  $h \circ f = h \circ g$ , then  $f = g$ .



**For next time:**

*Pg 346: 7.6.(2, 3, 6)*

*Ex "7.5.(a-c)" on Schoology*

*Pg 351: 7.8.(1, 5, 6)*

*Skim 7.9*