### Chapter 7 outline:


- Introduction, function equality, and anonymous functions (Monday)
- ► Image and inverse images (**Today**)
- ► Function properties, composition, and applications to programming (Friday)
- Cardinality (next week Monday)
- Countability (next week Wednesday)
- ► Review (Monday, Apr 18)
- ► Test 3, on Ch 6 & 7 (Wednesday, Apr 20)


## Today:

- Review definitions from last time
- New definitions: image and inverse image
- Proofs
- Programming



A relation f from X to Y is a function (written  $f: X \to Y$ ) if  $\forall x \in X$ , (1)  $\exists y \in Y \mid (x, y) \in f$ , and (2)  $\forall y_1, y_2 \in Y$ ,  $(x, y_1)$ ,  $(x, y_2) \in f \to y_1 = y_2$ .





Not a function.

(There's a domain element that is related to two things.)

Not a function.


(There's a domain element that is not related to anything.)

A function.

(It's OK that two domain elements are related to the same thing and one codomain element has nothing related to it.)


## **Image**

$$F(A) = \{ y \in Y \mid \exists \ x \in A \text{ such that } f(x) = y \}$$



## Inverse image

$$F^{-1}(B) = \{x \in X \mid f(x) \in B\}$$



**Lemma 7.2.** If  $f: X \to Y$ , then  $F(\emptyset) = \emptyset$ .


**Lemma 7.3.** If  $f: X \to Y$ ,  $A \subseteq X$ , and  $A \neq \emptyset$ , then  $F(A) \neq \emptyset$ .

**Lemma 7.4.** If  $f: X \to Y$ , then  $F^{-1}(\emptyset) = \emptyset$ .


We might expect the following, but it's not true:

**Lemma XXXX.** If  $f: X \to Y$ ,  $A \subseteq Y$ , and  $A \neq \emptyset$ , then  $F^{-1}(A) \neq \emptyset$ .

**Ex 7.4.1.** If  $A, B \subseteq X$ , then  $F(A \cap B) \subseteq F(A) \cap F(B)$ .



Consider this picture of X and Y:



**Attempted proof.** Suppose  $A, B \subseteq X$  and  $y \in F(A - B)$ . By definition of image, there exists  $x \in A - B$  such that f(x) = y.

**Attempted proof.** Suppose  $A, B \subseteq X$  and  $y \in F(A - B)$ . By definition of image, there exists  $x \in A - B$  such that f(x) = y.

By definition of difference,  $x \in A$ , and  $x \notin B$ . By definition of image,  $f(x) \in F(A)$ .

**Attempted proof.** Suppose  $A, B \subseteq X$  and  $y \in F(A - B)$ . By definition of image, there exists  $x \in A - B$  such that f(x) = y.


By definition of difference,  $x \in A$ , and  $x \notin B$ . By definition of image,  $f(x) \in F(A)$ . So, also by definition of image,  $f(x) \notin F(B)$ . Right?

**Attempted proof.** Suppose  $A, B \subseteq X$  and  $y \in F(A - B)$ . By definition of image, there exists  $x \in A - B$  such that f(x) = y.


By definition of difference,  $x \in A$ , and  $x \notin B$ . By definition of image,  $f(x) \in F(A)$ .

So, also by definition of image,  $f(x) \notin F(B)$ . Right?

NO!



**Ex 7.4.3.** If  $A, B \subseteq X$ , then  $F(A - B) \subseteq F(A) - F(B)$ ?



Let 
$$X = \{x_1, x_2\}$$
,  $Y = \{y\}$ ,  $A = \{x_1\}$ , and  $B = \{x_2\}$ .


Let 
$$f = \{(x_1, y), (x_2, y)\}.$$

Then 
$$F(A - B) = F(\{x_1\} - \{x_2\}) = F(\{x_1\}) = \{y\}.$$


Moreover, 
$$F(A) - F(B) = \{y\} - \{y\} = \emptyset$$
.

So 
$$F(A - B) \not\subseteq F(A) - F(B)$$


**Ex 7.4.4.** If  $A \subseteq B \subseteq X$ , then  $F(B) = F(B - A) \cup F(A)$ .



**Ex 7.4.6.** If  $A \subseteq B \subseteq Y$ , then  $F^{-1}(A) \subseteq F^{-1}(B)$ .



**Ex 7.4.7.** If  $A, B \subseteq Y$ , then  $F^{-1}(A \cup B) = F^{-1}(A) \cup F^{-1}(B)$ .



#### For next time:

Pg 342: 7.4.(2, 5, 8, 9, 10) (Programming problems are with the next assignment)

Read 7.(6-8)

Take quiz