

## Chapter 6 roadmap:


- ▶ Recursive definitions and types (last week Wednesday)
- ▶ Structural induction (Fridayday)
- ▶ Mathematical induction (**Today**)
- ▶ Loop invariant proofs (Wednesday and Friday)

Last time we saw self-referential proofs for propositions quantified over recursively defined sets, **structural induction**.

Today we see self-referential proofs for propositions quantified over the natural numbers and whole numbers.

- ▶ Opening examples and observations
- ▶ General form of **mathematical induction**
- ▶ Comments on the term *induction*
- ▶ Other examples, including on sets







Conjecture:

$$\forall n \in \mathbb{N}, \sum_{i=1}^n (2i - 1) = n^2$$

$$\sum_{i=1}^5 (2i - 1) = (2 \cdot 1 - 1) + (2 \cdot 2 - 1) + (2 \cdot 3 - 1) + (2 \cdot 4 - 1) + (2 \cdot 5 - 1) = 1 + 3 + 5 + 7 + 9$$

Recall the Peano definition of  $\mathbb{W}$ . Similarly for  $\mathbb{N}$ :  $n \in \mathbb{N}$  if  $n = 1$  or  $n = x + 1$  for some  $x \in \mathbb{N}$ .

$$\forall n \in \mathbb{N}, \sum_{i=1}^n (2i - 1) = n^2$$

$$\forall n \in \mathbb{N}, \sum_{i=1}^n (2i - 1) = n^2$$

**Proof.** Suppose  $n \in \mathbb{N}$ . Then either  $n = 1$  or there exists  $x \in \mathbb{N}$  such that  $n = x + 1$ .

**Base case.** Suppose  $n = 1$ . Then

$$\sum_{i=1}^1 (2i - 1) = 2 - 1 = 1 = 1^2$$

**Inductive case.** Suppose  $n = x + 1$  such that  $x \in \mathbb{N}$  and  $\sum_{i=1}^x (2i - 1) = x^2$ .  
Then

$$\begin{aligned} \sum_{i=1}^n (2i - 1) &= 2n - 1 + \sum_{i=1}^{n-1} (2i - 1) && \text{by definition of summation} \\ &= 2n - 1 + \sum_{i=1}^x (2i - 1) && \text{by substitution} \\ &= 2n - 1 + x^2 && \text{by the inductive hypothesis} \\ &= 2n - 1 + (n - 1)^2 && \text{by substitution} \\ &= 2n - 1 + n^2 - 2n + 1 && \text{by algebra (FOIL)} \\ &= n^2 && \text{by algebra (cancellation)} \quad \square \end{aligned}$$

$$4|0 \quad 0 + 1 = 1 = 5^0$$

$$4|4 \quad 4 + 1 = 5 = 5^1$$

$$4|24 \quad 24 + 1 = 25 = 5^2$$

$$4|124 \quad 124 + 1 = 125 = 5^3$$

$$4|624 \quad 624 + 1 = 625 = 5^4$$

Conjecture:  $\forall n \in \mathbb{W}, 4|5^n - 1$

$$\forall n \in \mathbb{W}, 4 \mid 5^n - 1$$



$$\forall n \in \mathbb{W}, 4|5^n - 1$$

**Proof.** By induction on  $n$ .

**Base case.** Suppose  $n = 0$ . Then  $5^0 - 1 = 1 - 1 = 0 = 4 \cdot 0$ . Hence  $4|5^0 - 1$  by the definition of divides.

**Inductive case.** Suppose  $n > 0$  and  $4|5^{n-1} - 1$ .

Then, by definition of divides, there exists  $k \in \mathbb{W}$  such that  $5^{n-1} - 1 = 4k$ .  
Moreover,

$$\begin{aligned} 5^n - 1 &= 5 \cdot 5^{n-1} - 1 && \text{by algebra, unless otherwise noted...} \\ &= 5 \cdot (5^{n-1} - 1 + 1) - 1 \\ &= 5(4k + 1) - 1 && \text{by the inductive hypothesis} \\ &= 5 \cdot 4 \cdot k + 5 - 1 \\ &= 5 \cdot 4 \cdot k + 4 \\ &= 4(5k + 1) \end{aligned}$$

Hence  $4|5^n - 1$  by definition of divides.  $\square$

$$\forall n \in \mathbb{W}, 4|5^n - 1$$

**Proof.** By induction on  $n$ .

**Base case.** Suppose  $n = 0$ . Then  $5^0 - 1 = 1 - 1 = 0 = 4 \cdot 0$ . Hence  $4|5^0 - 1$  by the definition of divides.

**Inductive case.** Suppose  $4|5^n - 1$  for some  $n \geq 0$ .

Then, by definition of divides, there exists  $k \in \mathbb{W}$  such that  $5^n - 1 = 4k$ .  
Moreover,

$$\begin{aligned} 5^{n+1} - 1 &= 5 \cdot 5^n - 1 && \text{by algebra, unless otherwise noted. . .} \\ &= 5 \cdot (5^n - 1 + 1) - 1 \\ &= 5(4k + 1) - 1 && \text{by the inductive hypothesis} \\ &= 5 \cdot 4 \cdot k + 5 - 1 \\ &= 5 \cdot 4 \cdot k + 4 \\ &= 4(5k + 1) \end{aligned}$$

Hence  $4|5^{n+1} - 1$  by definition of divides.  $\square$

To prove  $\forall n \in \mathbb{W}, I(n)$ ,

- ▶ Show  $I(0)$
- ▶ Show  $\forall n \in \mathbb{W}, I(n) \rightarrow I(n+1)$ , that is  
Suppose  $n \geq 0$  such that  $I(n)$

$\vdots$

$I(n+1)$

Alternately, show  $\forall n \in \mathbb{W}$  such that  $n > 0, I(n-1) \rightarrow I(n)$ , that is

Suppose  $n \geq 0$  such that  $I(n-1)$

$\vdots$

$I(n)$

- ▶ Conclude  $\forall n \in \mathbb{W}, I(n)$

The *principle of mathematical induction* is

$$[I(0) \wedge \forall n \in \mathbb{W}, I(n) \rightarrow I(n+1)] \rightarrow [\forall n \in \mathbb{W}, I(n)]$$

$$\sum_{i=1}^1 i = 1 = 1 = \frac{1 \cdot 2}{2}$$

$$\sum_{i=1}^2 i = 1 + 2 = 3 = \frac{2 \cdot 3}{2}$$

$$\sum_{i=1}^3 i = 1 + 2 + 3 = 6 = \frac{3 \cdot 4}{2}$$

$$\sum_{i=1}^4 i = 1 + 2 + 3 + 4 = 10 = \frac{4 \cdot 5}{2}$$

$$\sum_{i=1}^5 i = 1 + 2 + 3 + 4 + 5 = 15 = \frac{5 \cdot 6}{2}$$

**Ex 6.5.1.**  $\forall n \in \mathbb{N}, \sum_{i=1}^n i = \frac{n(n+1)}{2}$ .

**Ex 6.5.1.**  $\forall n \in \mathbb{N}, \sum_{i=1}^n i = \frac{n(n+1)}{2}$ .

**Proof.** By induction on  $n$ .

**Base case.** Suppose  $n = 1$ . Then  $\sum_{i=1}^1 i = 1 = \frac{1(1+1)}{2}$ .

**Inductive case.** Suppose that for some  $n \geq 1$ ,  $\sum_{i=1}^n i = \frac{n(n+1)}{2}$ . Then

$$\sum_{i=1}^{n+1} i = n + 1 + \sum_{i=1}^n i \quad \text{by definition of summation}$$

$$= n + 1 + \frac{n(n+1)}{2} \quad \text{by the inductive hypothesis}$$

$$= \frac{2n+2+n^2+n}{2} \quad \text{by algebra}$$

$$= \frac{n^2+3n+2}{2} \quad \text{"}$$

$$= \frac{(n+1)(n+2)}{2} \quad \text{"} \quad \square$$

Observe:

$$|A|$$

$$|\mathcal{P}(A)|$$

$$|\emptyset| = 0$$

$$|\{\emptyset\}| = 1$$

$$|\{a\}| = 1$$

$$|\{\emptyset, \{a\}\}| = 2$$

$$|\{a, b\}| = 2$$

$$|\{\emptyset, \{a\}, \{b\}, \{a, b\}\}| = 4$$

$$|\{a, b, c\}| = 3$$

$$|\{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}\}| = 8$$

Conjecture: For any finite set  $A$ ,  $|\mathcal{P}(A)| = 2^{|A|}$ .

**Theorem 6.5.** For all  $n \in \mathbb{W}$ , if  $A$  is a set such that  $|A| = n$ , then  $|P(A)| = 2^n$ .



**Theorem 6.5.** For all  $n \in \mathbb{W}$ , if  $A$  is a set such that  $|A| = n$ , then  $|\mathcal{P}(A)| = 2^n$ .

**Proof.** By induction on  $n$ .

**Base case.** Suppose  $n = 0$ . Then  $A = \emptyset$ , and  $|\mathcal{P}(A)| = |\{\emptyset\}| = 1 = 2^0$ .

**Inductive case.** Suppose for some  $n \geq 0$ , if  $A$  is a set such that  $|A| = n$ , then  $|\mathcal{P}(A)| = 2^n$ . Suppose further that  $A$  is a set such that  $|A| = n + 1$ .

Since  $|A| > 0$ , let  $a \in A$ . By Corollary 4.12,  $\mathcal{P}(A - \{a\})$  and  $\{C \cup \{a\} \mid C \in \mathcal{P}(A - \{a\})\}$  make a partition of  $\mathcal{P}(A)$ . Then

$$\begin{aligned} |\mathcal{P}(A - \{a\})| &= |\{C \cup \{a\} \mid C \in \mathcal{P}(A - \{a\})\}| && \text{by Exercise 7.9.6} \\ |A - \{a\}| &= |A| - |\{a\}| && \text{since } \{a\} \subseteq A, \text{ and by Ex 7.9.1} \\ &= n + 1 - 1 && \text{by supposition} \\ &= n && \text{by arithmetic} \\ |\mathcal{P}(A - \{a\})| &= 2^n && \text{by the inductive hypothesis} \\ |\mathcal{P}(A)| &= |\mathcal{P}(A - \{a\})| \\ &\quad + |\{C \cup \{a\} \mid C \in \mathcal{P}(A - \{a\})\}| && \text{by Theorem 7.12} \\ &= 2^n + 2^n && \text{by substitution} \\ &= 2^{n+1} && \text{by algebra. } \square \end{aligned}$$

Iterated union (similar for intersection):

$$\bigcup_{i=1}^n A_i = A_1 \cup A_2 \cup \cdots \cup A_n$$

**Ex 6.6.1.**  $\forall n \in \mathbb{N}, \overline{\bigcup_{i=1}^n A_i} = \bigcap_{i=1}^n \overline{A_i}$

**Proof.** *By induction on  $n$ .*

**Base case.** *Suppose  $n = 1$ . Then*

$$\overline{\bigcup_{i=1}^1 A_i} = \overline{A_1} = \bigcap_{i=1}^1 \overline{A_1}$$

**Inductive case.** Suppose  $\overline{\bigcup_{i=1}^n A_i} = \bigcap_{i=1}^n \overline{A_i}$  for some  $n \geq 1$ . Then

$$\begin{aligned}\overline{\bigcup_{i=1}^{n+1} A_i} &= \overline{A_{n+1} \cup \bigcup_{i=1}^n A_i} \quad \text{by definition of iterated union} \\ &= \overline{A_{n+1}} \cap \overline{\bigcup_{i=1}^n A_i} \quad \text{by Ex 4.3.13 (DeMorgan's law of sets)} \\ &= \overline{A_{n+1}} \cap \bigcap_{i=1}^n \overline{A_i} \quad \text{by the inductive hypothesis} \\ &= \bigcap_{i=1}^{n+1} \overline{A_i} \quad \text{by the definition of iterated intersection}\end{aligned}$$

□

**For next time:**

*Pg 273: 6.5.(2 & 4)*

*Pg 278: 6.6.(2 & 3)*

*Read 6.9 carefully*

*Skim 6.10*

*Take quiz (by Friday)*